SIGNS OF INFECTION & SEPSIS

Surgery can put you at increased risk for developing infection and sepsis. After surgery, patients and caregivers should watch for the following signs and symptoms.		
GREEN ZONE	 My heartbeat is as usual. Breathing is normal for me I have not had a fever in the past 24 hours and I am not taking medicine for a fever I do not feel chilled My energy level is as usual My thinking is clear I feel well I have taken my antibiotics as prescribed My wound or IV site is not painful, red, draining pus or smelling bad 	DOING GREAT! No action is needed.
YELLOW ZONE	 My heartbeat is faster than usual My breathing is a bit more difficult and faster than usual I have a fever between 100.5°F to 101.2°F I feel chilled and cannot get warm. I am shivering or my teeth are chattering I am too tired to do most of my usual activities I feel confused or I am not thinking clearly I do not feel well I have a bad cough or my cough has changed How often I urinate has changed. When I do urinate, it burns, is cloudy or smells bad My wound or IV site has changed 	TAKE ACTION TODAY! Call your doctor's office. (Phone number)
RED ZONE	 My heartbeat is very fast My breathing is very fast and more difficult My temperature is below 96.8°F. My skin or fingernails are pale or blue My fever is 101.2°F or more I have not urinated for 5 or more hours I am very tired. I cannot do any of my usual activities My caregivers tell me I am not making sense I feel sick My cough is much worse 	TAKE ACTION NOW! Go to the Emergency Room AND Call your doctor's office. (Phone number)

Sources: Sepsis Alliance, sepsis.org; Centers for Disease Control and Prevention (CDC), cdc.gov; and atom Alliance, atomalliance.org

• My wound or IV site is painful, red, smells bad or has pus

