

Cold Knife Conization

Technique

Procedure involves the removal of the portion of your cervix that is involved with the dysplasia (precancerous cells)

Incisions

A small incision is made on the cervix (mouth of the womb).

Operative Time

Operative times vary greatly depending on the findings at the time of the procedure. Your surgeon will proceed with safety as his/her priority. Average times range from 15-30 minutes.

Anesthesia

- Local anesthesia + IV sedation
- General anesthesia

Preoperative Care

- Schedule your case immediately after your period.
- Nothing by mouth after midnight

Hospital Stay

Day surgery

Postoperative Care

These guidelines are intended to give you a general idea of your postoperative course.

Since every patient is unique and has a unique procedure, your recovery may differ.

- An-inflammatory pain medicine, such as ibuprofen, naproxen, etc., is usually required for the first several days.
- Driving is allowed on the day after your procedure.
- If they desire, patients may return to work on the day after the procedure.
- You may experience mild vaginal bleeding for several days after the procedure. If it is not associated with increasing pain or fever, it is nothing to worry about and will resolve.

Physical Address: 1700 6th Ave. S.

Mailing Address: 176F, Ste. 9103, 619 19th St. S., Birmingham, AL 35249

Phone: 205-996-3130 | Fax: 205-996-3170